

Annex 1: Data Collection Tool

Questionnaire

Patients' information

1. Age (years) :
2. Hospital settings Al-Shifa Hospita Nasser medical complex
3. BMI?
4. Parity: Nulliparous Multiparous: number of births
5. Antenatal Care : Yes No
6. Previous history of PPH : Yes If yes how many times?..... No
7. Previous history of Caesarean section : Yes No
8. Previous history of Gynecological operations : Yes No
9. Previous history of Rupture of the uterus in childbirth : Yes No
10. Previous history of multiple gestation: Yes No
10. Previous history of multiple gestation: Yes No
11. Baby weight : <4000 g >4000 g
12. History of APH in this current pregnancy? Yes No
13. Placental site: normally-sited, Low lying, previa
14. was this pregnancy: induced? spont?
15. Birth weight of baby?.....
16. delivery: term, post-term , pre-term
17. current history of chorioamnionitis? Yes No
18. history of polyhy..? Yes No
19. History of intrahepatic cholestasis? Yes No
20. History of coagulation diseases? Yes No

21. any anticoagulant? Yes No
if yes: aspirin, clexan , when it was stopped?.....
22. anemia Yes No

Prevention of Postpartum Hemorrhage (PPH) : Active management of III stage of labor:

1. Give Oxytocin 10 unit IM or IV injection with delivery of the anterior shoulder or immediately after delivery for the fetus

Yes No

if not: what is the dose and route?.....

2. Delay cord clamping for at least 1-3 minutes

Yes No

3. Deliver the placenta using Controlled Cord Traction (CCT)

Yes No

4. Perform uterine massage to the fundus of the uterus through the woman's abdomen until the uterus is contracted

Yes No

5. Repeat uterine massage every 15 minutes for the first hour

Yes No

6. Document all procedures, medications given, assessment findings

Yes No

For high risk cases :

1. Give Methergine IM (1 ml : 0.25 mg) After delivery and exam to confirm complete placental delivery?

Yes No

2. Oxytocin 40 units in 1000 cc NS at a rate of 250 mm/hour and/or Misoprostol 800 – 1000 mcg PR

Yes No

3. are these measures done routinely as prophylaxis to PPH according to national protocol?

Yes No