

# Beyond PaO<sub>2</sub>: Relationships between Indices of Pulmonary Oxygen Transfer in Hospitalized Adults Undergoing Blood Gas Testing

Mark Feldman, MD, Denise Rebel, BS, RRT

Department of Internal Medicine and the Pulmonary Lab, Texas Health Presbyterian Hospital Dallas, Dallas, TX, USA

\*Author of Correspondence: Mark Feldman, MD; [markfeldman@texashealth.org](mailto:markfeldman@texashealth.org)

Received 24 February 2019;

Accepted 12 March 2019;

Published 14 March 2019

## Abstract

The arterial partial pressure of oxygen in the arterial blood reveals limited information about pulmonary oxygen transfer. In 989 hospitalized patients undergoing arterial blood gas testing, we compared conventional indices of oxygen transfer: (1) P(A-a) O<sub>2</sub> (A-a); (2) PaO<sub>2</sub>/PAO<sub>2</sub> (a/A); and (3) the P/F ratio. Nine hundred twenty-five of the patients were receiving supplemental oxygen therapy (FIO<sub>2</sub> .24-1.00) and 65 were breathing room air. In patients receiving supplemental O<sub>2</sub>, the a/A ratio closely correlated with the P/F ratio ( $r = .98$  to  $.99$ ); the A-a O<sub>2</sub> difference did not correlate as closely with the a/A ratio ( $r = -.77$  to  $-.85$ ) or with the P/F ratio ( $r = -.72$  to  $-.80$ ). In those breathing room air (FIO<sub>2</sub> .21), the a/A ratio was closely and inversely correlated with the A-a difference ( $r = -.97$  to  $-.98$ ); correlations between the a/A and P/F ratios ( $r = .88$  to  $.89$ ) and between the P/F ratio and A-a O<sub>2</sub> difference ( $r = -.78$  to  $-.79$ ) were less robust. We conclude that the a/A ratio is the preferred oxygen transfer parameter over the wide range of FIO<sub>2</sub> levels encountered clinically.

**Keywords:** alveolar oxygen pressure, arterial oxygen pressure, blood gases, hypoxemia, lung oxygen transfer, oxygen, carbon dioxide.

## Introduction

The PaO<sub>2</sub> is sometimes used to determine how well the lungs are transferring oxygen to the circulation. Unfortunately, an isolated PaO<sub>2</sub> measurement conveys limited O<sub>2</sub> exchange information unless PAO<sub>2</sub> is also considered.<sup>[1]</sup> A larger-than-expected difference between the PAO<sub>2</sub> and the PaO<sub>2</sub>, the A-a O<sub>2</sub> difference, provides a reliable indication of an abnormality of pulmonary O<sub>2</sub> transfer.<sup>[1,2]</sup> A larger-than-expected A-a difference can be caused by a pulmonary ventilation-perfusion imbalance, a barrier to O<sub>2</sub> diffusion across the alveoli, shunting of some deoxygenated venous blood through or past the lungs, or some combination of these factors.<sup>[1]</sup>

PAO<sub>2</sub> is directly related to the alveolar ventilation rate, increasing when there is alveolar hyperventilation and decreasing when there is alveolar hypoventilation. Calculation of PAO<sub>2</sub> (to determine A-a O<sub>2</sub>) is usually made by using the alveolar gas equation.<sup>[2]</sup> Use of this time-honored equation relies on numerous assumptions: (a) estimated FIO<sub>2</sub> is accurately known; (b) ambient P<sub>b</sub> is precisely known; (c) PICO<sub>2</sub> is 0; (d) PaCO<sub>2</sub> equals PACO<sub>2</sub>; (e) the water vapor pressure in the trachea and pulmonary alveoli is 47 mmHg; and (f) the respiratory exchange ratio is 0.8.<sup>[1,2]</sup> The A-a O<sub>2</sub> difference increases with aging and also with use of supplemental

O<sub>2</sub> therapy, and well-defined normal ranges for A-a are not known at all possible age and FIO<sub>2</sub> combinations.<sup>[1]</sup>

The a/A ratio is an improvement on the A-a because it is unaffected by the FIO<sub>2</sub>. Thus, a/A can be assessed serially even when the FIO<sub>2</sub> has changed over time. The a/A ratio may be the preferred index for quantitating O<sub>2</sub> transfer by the lung.<sup>[3]</sup> However, a/A remains dependent on the validity of the numerous assumptions made to calculate PAO<sub>2</sub>. The a/A ratio is also age-related, being  $>0.75$  in healthy older individuals and often  $>0.9$  in younger people.

The P/F ratio is the simplest of the pulmonary O<sub>2</sub> transfer indices and only requires an accurate estimate of FIO<sub>2</sub>. The P/F ratio calculation does not take into consideration the ambient P<sub>b</sub> or the alveolar ventilation rate as reflected by the patient's PaCO<sub>2</sub>. If the ambient P<sub>b</sub> is typical for the locale and the PaCO<sub>2</sub> is close to normal, the P/F should be a very accurate reflection of O<sub>2</sub> transfer by the lung. However, fluctuations in P<sub>b</sub> or PaCO<sub>2</sub> from typical or normal levels occur not uncommonly and can introduce inaccuracy if P/F is used as an index of lung oxygen transport. P/F can be estimated noninvasively by measuring using SO<sub>2</sub> (%) and using tables that convert SO<sub>2</sub>(%) to estimated PaO<sub>2</sub>.<sup>[4]</sup> However, these tables require assumptions about arterial pH, body temperature, and erythrocyte 2,3-DPG levels.

Often, A-a, a/A, and P/F are reported by the laboratory to the care provider who ordered the blood gas measurement. The amount of detail provided may be confusing and redundant for a busy clinician. To what extent A-a, a/A, and P/F are interrelated in hospitalized patients undergoing blood gas measurements is unclear. We therefore analyzed 1,000 consecutive arterial blood gas patient samples that had been collected in our hospital and analyzed in our laboratory. Most patients were receiving supplemental oxygen therapy. A smaller subgroup breathing room air was included for comparison.

**Methods**

One thousand consecutive arterial blood gas samples collected in our hospital during March and April 2008 were evaluated for inclusion in the study. All samples had been obtained from adults (age 18 and above). To maintain patient confidentiality, samples were de-identified in the laboratory by assigning them a sequential study number from 1 to 1,000 before sending them to the investigator analyzing the data. Most samples were collected from patients who were receiving supplemental oxygen therapy (Table 1).

**Table 1: FIO2 recorded at the time of collection of the arterial blood gas samples<sup>a</sup>**

	FIO2	Number of patients/samples
Breathing Room Air (n=65)	.21	65
Receiving Supplemental O2 (n=924)	.24-.29	32
	.3-.39	80
	.4-.49	384
	.5-.59	73
	.6-.69	74
	.7-.79	95
	.8-.89	33
	.9-.99	9
	1	144

<sup>a</sup>11 of the original 1,000 blood gas samples (1.1%) were excluded due to quality control measures (see text). To calculate PIO2 in mmHg from FIO2, multiply FIO2 by (Pb-47).

The Abbott iStat<sup>®</sup> blood gas analyzer (Arbroath Angus, UK) was used to measure arterial PaO2, PaCO2, and pH in the samples. From these analytes and from the estimated FIO2 recorded at the time of sample collection, the following were calculated: (a) arterial SO2(%) from PaO2 using the Severinghaus formula;<sup>[5]</sup> (b) arterial HCO3 and base excess concentrations (mM) from the PaCO2 and pH using the Henderson-Hasselbalch and the Siggaard-Anderson equations,<sup>[6]</sup> respectively; and (c) PAO2 (mmHg) using the alveolar gas equation, which inputs PaCO2, Pb, and PIO2<sup>[1,2]</sup> Lastly, the laboratory calculated and reported three conventional oxygenation indices: A-a O2 difference (mmHg); a/A ratio (no units); and P/F ratio (mmHg).

Eleven of the 1,000 arterial samples (1.1%) were excluded if there was a negative A-a O2 difference (a/A ratio >1), implying an error in recording the true FIO2, or if the FIO2 was recorded as <0.21 or >1.00. The remaining 989 samples/patients were included for analysis.

Paired correlations of A-a, a/A, and P/F were calculated using the Pearson method (Excel, Microsoft Office Professional Plus 2013, Redmond, WA) and the Spearman rank-order correlation

method.<sup>[6]</sup> Linear regression analysis was performed to determine slopes and y-intercepts.<sup>[7]</sup> Group means (supplemental O2 vs. room air groups) were compared by unpaired t-tests, with 2-tailed P values < 0.05 considered significant.<sup>[8]</sup>

**Results**

Table 2 shows the laboratory’s reference range for the arterial analytes (PaO2, PaCO2, and pH) and the derived values of SO2(%), HCO3, and base excess. There were highly significant differences between mean values in the two groups of patients for all six parameters. Compared to patients breathing supplemental O2, those breathing room air had lower mean values for all six parameters.

**Table 2: Comparison of arterial PaO2, PaCO2 and pH as well as derived SO2(%),HCO3 level and base excess levels in the two groups of patients**

Parameter (RR)	Supplemental O2 (n=924)	Room Air (n=65)	P value
	Mean ±SEM (Range)	Mean ±SEM (Range)	
PaO2 (mmHg) (80-105 <sup>a</sup> )	134±3 (27-623)	78±3 (26-121)	<0.001
PaCO2 (mmHg) (35-45)	40±.4 (15-107)	36±1 (18-62)	0.009
pH (7.35-7.45)	7.40±.003 (6.88-7.68)	7.36±.01 (7.17-7.51)	0.006
SO2 (%) (96-100)	97±.1 (40-100)	92±2 (20-99)	<0.001
HCO3 (mM) (22-26)	24.7±.2 (4.6-50.9)	21.0±.9 (6.9-40.2)	<0.001
Base excess (mM) (-2-+3)	-0.1±.2 (-26 - +30)	-4.4±1.1 (-22 - +16)	<0.001

<sup>a</sup>based on breathing room air

Table 3 shows mean A-a O2 differences, a/A ratios, and P/F ratios in the two groups of patients.

**Table 3: Calculated A-a, a/A and P/F in the two groups of patients**

Index (RR)	Supplemental O2 (n=924)	Room Air (n=65)	P value
	Mean± SEM (Range)	Mean± SEM (Range)	
A-a (mm Hg) (4-25 <sup>a</sup> )	216±5 (0-644)	27±2 (0-80)	<0.001
a/A (≥0.75)	0.42±.007 (0.04-1.00)	0.74±.02 (0.25-1.00)	<0.001
P/F (mm Hg) (380-500)	257±49 (27-263)	369±12 (124-576)	<0.001

F, FIO2 expressed as a decimal; P, PaO2 in mmHg

<sup>a</sup>based on breathing room air


The supplemental O2 group (FIO2 .24-1.00), as expected, had markedly impaired O2 transfer as reflected by each of the oxygenation parameters (Table 3). In this group, there was a very close direct correlation (r= .98 and .99) between the a/A ratio and the P/F ratio (Table 4 and Figure 1A).

**Table 4: Correlation coefficients (r) between pairs of oxygenation indices<sup>a</sup>**

Supplemental O2 Group (n=924)			
Pearson (parametric)		Spearman (non-parametric)	
a/A vs. P/F	r= .98	r= .99	a/A vs. P/F
a/A vs. A-a	r= -.77	r= -.85	a/A vs. A-a
P/F vs. A-a	r= -.72	r= -.80	P/F vs. A-a
Room Air Group (n=65)			
a/A vs. P/F	r= .89	r= .88	a/A vs. P/F
a/A vs. A-a	r= -.98	r= -.97	a/A vs. A-a
P/F vs. A-a	r= -.78	r= -.79	P/F vs. A-a


<sup>a</sup>The r values can vary from -1 to +1. All r values reported here are significant (P<0.001).

The linear regression equation relating a/A to P/F was:  $(a/A)=.0017(P/F)+.0027$ . The 95% confidence interval for the .0017 slope of this relationship was .0016 to .0017. Thus, for every 100-mmHg increase (or decrease) in P/F, a/A increased (or decreased) by 0.16-0.17. In this supplemental O2 group, there were less robust inverse correlations between a/A and A-a (Table 4 and Figure 1B) and between P/F and A-a (Table 4).



In patients breathing room air, the indices of oxygenation were close to the reference ranges (Table 3), implying that the blood gas sample had been ordered more to assess pH or PaCO<sub>2</sub> than PaO<sub>2</sub> (see also Table 2). In this group there was a very close inverse correlation between a/A and A-a (r = -.98 and -.97) (Table 4 and Figure 2A). The linear regression equation relating them was:

$(a/A)=0.984-.009(A-a)$ . The 95% confidence interval for the -.009 slope of this relationship was -.0086 to -.0096. Thus, for every 100 mmHg increase in A-a, a/A would fall by approximately 0.09. There were more modest correlations between a/A and P/F (Table 4 and Figure 2B) and (inversely) between P/F and A-a in this group (Table 4).



## Discussion

Results of this study varied depending on the group of hospitalized patients analyzed. In the larger group receiving a wide range of supplemental O<sub>2</sub> doses (FIO<sub>2</sub> 0.24-1.00), a/A correlated very closely with P/F, but not nearly as well with A-a. This finding was not entirely unexpected, as a/A is independent of FIO<sub>2</sub>, whereas the A-a O<sub>2</sub> difference is highly FIO<sub>2</sub>-dependent. The close correlation between a/A and P/F implies that the parameters used to calculate PAO<sub>2</sub> are not as critical as assessing oxygen transfer by the lungs in patients receiving supplemental O<sub>2</sub> as they might be in people breathing room air. These parameters include the ambient P<sub>b</sub> and the patient's PaCO<sub>2</sub>. The 0.0017 slope of the line relating P/F to a/A with its narrow confidence interval and its y-intercept close to zero (Figure 1A) indicate that a/A can be easily and closely estimated from P/F and vice versa for inpatients receiving O<sub>2</sub> therapy.

A smaller group of patients underwent arterial blood gas testing while breathing room air, many of whom had a metabolic acidosis with respiratory compensation (Table 2). In this group, the a/A ratio better correlated with A-a than with P/F. However, in this

group FIO<sub>2</sub> was constant (0.21), unlike in the supplemental O<sub>2</sub> group where FIO<sub>2</sub> varied >4-fold. Thus, in hospitalized patients breathing room air, a/A and A-a (which use the same parameters) are highly inversely correlated and give essentially identical information. The -0.009 slope of the line relating A-a and a/A, with its narrow confidence interval and its y-intercept approaching 1, indicate that a/A can be easily and closely estimated from A-a and vice versa. In contrast to the supplemental O<sub>2</sub> group, the a/A ratio in the room air group was not as highly correlated with the P/F ratio, probably because inclusion of the ambient P<sub>b</sub> and PaCO<sub>2</sub> in the a/A calculation, but not the P/F calculation, provided less precise information for P/F than for a/A in this group of subjects.

## Conclusion

We agree with an editorial opinion expressed nearly 35 years ago by Hess and Maxwell that the a/A ratio is the preferred parameter of oxygenation and lung O<sub>2</sub> transfer whether patients are breathing room air or receiving supplemental O<sub>2</sub>. For the former group, the A-a O<sub>2</sub> difference is a suitable alternative to a/A, while for the latter group, the P/F ratio provides a suitable alternative to a/A. Another advantage of relying on the a/A ratio is that it has no units.

Both A-a and P/F can be expressed in different pressure units globally (e.g., mmHg, torr, or kilopascals), which can lead to confusion when comparing results across institutions or groups of investigators. Our conclusions apply to patients admitted to the hospital in whom arterial blood gases were ordered by their care provider regardless of diagnosis. For certain known diagnoses (e.g., pneumonia, pulmonary embolism, acute respiratory distress syndrome), PaO<sub>2</sub> alone, the P/F ratio or the A-a O<sub>2</sub> difference might perform equally or even superiorly to the a/A ratio. Additional studies may shed further light on this issue.

## References

- [1] Martin L. Abbreviating the alveolar gas equation: An argument for simplicity. *Respiratory Care* 1985; 30: 964-8.
- [2] Fenn WO, Rahn H, Otis AB. A theoretical study of the composition of the alveolar air at altitude. *Amer J Physiol* 1946; 146: 637-653.
- [3] Hess D, Maxwell C. Editorial. Which is the best index of oxygenation- P(A-a)O<sub>2</sub>, PaO<sub>2</sub>/PAO<sub>2</sub>, or PaO<sub>2</sub>/FIO<sub>2</sub>? *Respiratory Care* 1985; 30: 961-3.
- [4] Conversion tables. <https://www.intensive.org/Documents>. Last retrieved 02/15/19.
- [5] Severinghaus JW. Simple, accurate equations for human blood O<sub>2</sub> dissociation computations. *J Appl Physiol* 1979; 46: 599-602.
- [6] Siggaard-Anderson O, Fogh-Andersen N. Base excess or buffer base (strong ion difference) as a measure of non-respiratory acid-base disturbance. *Acta Anaesthesiol Scand Suppl* 1995; 107: 123-8.
- [7] Correlation and Regression. VassarStats: Statistical Web Site. <http://vassarstats.net>. Last retrieved 02/15/19.
- [8] GraphPad Software. [www.graphpad.com](http://www.graphpad.com). Last retrieved 02/16/19.

## Figure Legends

- 1A. Direct and generally close linear relationship between a/A and P/F in 924 patients receiving supplemental O<sub>2</sub>. Correlation coefficients were .98 (Pearson) and .99 (Spearman) (Table 4). Linear regression equation was: (a/A)= 0.0017(P/F) + 0.0027. The 95% confidence interval for the slope was 0.0016 to 0.0017. 1B. Nonlinear relationship between a/A and A-a in these same patients. Correlation coefficients were only -.77 and -.85 (Table 4). Trendlines are shown.
- 2A. Inverse linear relationship between a/A and A-a in 65 patients breathing room air. Correlation coefficients were -.98 (Pearson) and -.97 (Spearman) (Table 4). Linear regression equation was: (a/A)= 0.984-0.009(A-a). The 95% confidence interval for the slope was - 0.0086 to - 0.0096. 2B. Direct, moderate linear relationship between a/A and P/F in the same patients. Correlation coefficients were only .88 and .89 (Table 4). Trendlines are shown.

## Abbreviations

A-a, PAO<sub>2</sub>-PaO<sub>2</sub>  
a/A, PaO<sub>2</sub>/PAO<sub>2</sub>  
2,3-DPG, 2,3-diphosphoglycerate  
FIO<sub>2</sub>, fraction of inspired oxygen  
HCO<sub>3</sub>, bicarbonate  
PACO<sub>2</sub>, alveolar carbon dioxide pressure  
PaCO<sub>2</sub>, arterial carbon dioxide pressure  
PAO<sub>2</sub>, alveolar oxygen pressure  
PaO<sub>2</sub>, arterial oxygen pressure  
Pb, barometric pressure  
P/F, PaO<sub>2</sub>/FIO<sub>2</sub>  
PICO<sub>2</sub>, inspired carbon dioxide pressure  
PIO<sub>2</sub>, inspired oxygen pressure  
RR, reference range  
SEM, standard error of the mean  
SO<sub>2</sub>(%), percentage oxygen saturation of hemoglobin